


**NYSTATIN- nystatin suspension**  
**VistaPharm, Inc.**

-----  
**NYSTATIN ORAL SUSPENSION, USP 100,000 Units/mL**

**DESCRIPTION**

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei*. Structural formula:



Nystatin Oral Suspension, for oral administration, contains 100,000 USP Nystatin Units per mL. Inactive ingredients: alcohol ( $\leq 1\%$  v/v), methylparaben, NF; dibasic sodium phosphate, USP; monobasic sodium phosphate, USP; saccharin sodium, USP; sucrose (50% w/v), NF; glycerin, USP; carboxymethylcellulose sodium, USP; propylparaben, NF; artificial wild cherry flavor # 14783 and purified water, USP

**CLINICAL PHARMACOLOGY**

**Pharmacokinetics**

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

**Microbiology**

Nystatin is both fungistatic and fungicidal in vitro against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin in vitro on repeated subculture in increasing levels of nystatin; other *Candida* species become quite resistant. Generally, resistance does not develop in vivo. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intra-cellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

**INDICATIONS AND USAGE**

Nystatin Oral Suspension is indicated for the treatment of candidiasis in the oral cavity.

## CONTRAINDICATIONS

The preparation is contraindicated in patients with a history of hypersensitivity to any of its components.

## PRECAUTIONS

### General

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

### Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

### Pregnancy

#### Teratogenic Effects

##### Category C

Animal reproduction studies have not been conducted with nystatin oral suspension. It is also not known whether nystatin oral suspension can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin oral suspension should be given to a pregnant woman only if clearly needed.

### Nursing Mothers

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

### Pediatric Use

See **DOSAGE AND ADMINISTRATION**.

## ADVERSE REACTIONS

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General**).

*Gastrointestinal:* Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

*Dermatologic:* Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

*Other:* Tachycardia, broncho-spasm, facial swelling, and non-specific myalgia have also been rarely reported.

## OVERDOSAGE

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (see **CLINICAL PHARMACOLOGY, Pharmacokinetics**).

## DOSAGE AND ADMINISTRATION

**INFANTS:** 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place

one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

NOTE: Limited clinical studies in premature and low birth weight infants indicate that 1 mL four times daily is effective.

CHILDREN AND ADULTS: 4-6 mL (400,000 to 600,000 units) four times daily (one-half of dose in each side of mouth). The preparation should be retained in the mouth as long as possible before swallowing.

Continue treatment for at least 48 hours after perioral symptoms have disappeared and cultures demonstrate eradication of *Candida albicans*.

## **HOW SUPPLIED**

Nystatin Oral Suspension, USP, 100,000 USP Nystatin Units per mL, is available as a cherry flavored, light creamy yellow, ready-to-use suspension. It is supplied as follows:

NDC 66689-037-01. 5 mL unit dose cup.

NDC 66689-037-50. Case contains 50 unit dose cups of 5 mL (NDC 66689-037-01), packaged in 5 trays of 10 unit dose cups each.

NDC 66689-037-99. Case contains 100 unit dose cups of 5 mL (NDC 66689-037-01), packaged in 10 trays of 10 unit dose cups each.

## **Storage**

Store at 20°-25°C (68°-77°F) [See USP Controlled Room Temperature]. Avoid freezing.

## **Rx Only**

### **Manufactured by:**

VistaPharm, Inc.  
Largo, FL 33771

VP2053R1  
02/17

## **PRINCIPAL DISPLAY PANEL**

**Xact DOSE™**

**NYSTATIN**

**ORAL SUSPENSION, USP**

**500,000 units/ 5 mL**

NDC 66689-037-01

**Alcohol not more than 1% v/v**

**SHAKE WELL BEFORE USING**

**Protect from freezing**

**Store at 20°-25°C (68°-77°F)**

[See USP CONTROLLED RM TEMP]

**Rx Only This cup delivers 5 mL VP2052R1**

Manufactured by VistaPharm, Inc.  
Largo, FL 33771



**NYSTATIN**  
**ORAL SUSPENSION, USP**  
**500,000 units/5 mL**

NDC 66689-037-01



Alcohol not more than 1% v/v  
**SHAKE WELL BEFORE USING**  
Protect from freezing

Store at 20°-25°C (68°-77°F)  
[See USP CONTROLLED RM TEMP]

**Rx Only** This cup delivers 5 mL VP2052R1  
Manufactured by VistaPharm, Inc.  
Largo, FL 33771

## NYSTATIN

nystatin suspension

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66689-037
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E)	NYSTATIN	100000 [USPU] in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
METHYL PARABEN (UNII: A2I8C7HI9T)	
SODIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: 22ADO53M6F)	
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
GLYCERIN (UNII: PDC6A3C0OX)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
ALCOHOL (UNII: 3K9958V90M)	

PROPYLPARABEN (UNII: Z8IX2SC1OH)

### Product Characteristics

<b>Color</b>	YELLOW (Light yellow)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66689-037-50	5 in 1 CASE	05/10/2010	
1		10 in 1 TRAY		
1	NDC:66689-037-01	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		
2	NDC:66689-037-99	10 in 1 CASE	05/10/2010	
2		10 in 1 TRAY		
2	NDC:66689-037-01	5 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA064142	05/10/2010	

**Labeler** - VistaPharm, Inc. (116743084)

Revised: 7/2017

VistaPharm, Inc.